

STRATIGRAPHIC LEGEND

QUATERNARY

SOUTH-EAST CHURCHILL PROVINCE

TORNGAT OROGEN

Hutton anorthosite Complex: recrystalized anorthosite gneiss, granoblastic

Quartz "mesocrate" diorite, gray to black weathering. Deformed, foliated on outer edge of the Abloviak Shear zone; contains enclaves of porphoritic igneous rocks defined by aggregates of recrystalized plagioclase, and with <50 cm syngenetic xenoliths of fine granulitic diorite; transformed to mylonite in the Abloviak Shear

Mafic gneiss and amphibolite, characterized by compositional units having a centimetre thickness , made of amphibolite with plagioclase phenocrysts and other amphibolite units with a homogenous appearance.

Marble impure calcsilicate, consisting of 1-5 m beds of impure marble with diopside-olivine-calcite, and pale green 1-10 m thick diopside calc-silicate beds, locally numerous stratiform and discordant white quartz veins.

Paragneiss and migmatite metasediments, gray to tan weathering, compositionally characterized by red garnets, biotite and abundant leucosome in which sillimanite

Paragneiss and "diatexite", rusty brown weathering and white weathering, characterized by lilac garnets and sillimanite. Homogenous on a large scale, but well layered on a scale of 10cm-2m varies from a quartzite to garnet to semipelitic gneiss to garnet-biotite-feldspar-quartz, to quartzfeldspathic pelitic gneiss to

BURWELL DOMAIN PALEOPROTEROZOIC

Polyphasic intrusive suite, composed of diorite, tonalite, granodiorite, granite and quartziferous monzodiorite, metamorphosed to amphibolite facies, to granulite and retrogressive granulitic assemblages. This unit includes homogenous leucotonalite to quartz diorite, usually foliated (>80%), with inclusions of diorite and gabbro with plagioclase phenocrysts and stratified mafic gneiss, this rock is cut by numerous felsic and pegmatite veins.

Killinek Charnockite Suite
Granodiorite to homogenous tonalite with orthopyroxene (opdalite to enderbite),
tan weathering massive to foliated, equigranular medium grained rock.

Medium grained mafic diorite, having genoblastic texture, often found at the margins of charnockites (pPkil2) and forming flakes in paleoproterozoic age

ARCHEAN to PALEOPROTEROZOIC

having been affected and deformed during the Torngat Orogen, having undergone retrograde metamorphism generally to upper amphibolite facies in the paleoproterozoic, locally some granulite metamorphic facies mineral assemblages

NORTH CRATON **PALEOPROTEROZOIC**

Biotite Paragneiss, sillimanite garnet with brown weathering alteration.

Calcitic marble, dolomitic marble, zones of calcsilicate with diopsides and local

Biotite paragneiss, quartzous paragneiss and garnet biotite paragneiss with local

ARCHEAN to PALEOPROTEROZOIC

ApP kan1 White and grey tonalitic gneiss and migmatite with beds of homogenous mafic

LITHOLOGY LEGEND

BURWELL DOMAIN PALEOPROTEROZOIC

Fault breccia and cataclastite with associated veins of pseudotachylite.


Mylonite and ultramylonite at amphibolite and greenschist facies.


Fine to medium grained pink to gray granite having a sugary texture.


MRNFP-GÉOINFORMATION 2004


GM 60874


Reference: Natural Resources Quebec, 2001 1:50 000 Geology map sheet 24P07-MONT JACQUES-ROUSSEAU SI-24 P07-C3G-01H, Van Kranendonk, M.J.























GM 60874

